

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrKALYDECO®

Ivacaftor Tablets Ivacaftor Granules

Read this carefully before you/your child start taking **KALYDECO** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your/your child's medical condition and treatment and ask if there is any new information about **KALYDECO**.

What is KALYDECO used for?

KALYDECO tablets (150 mg) are for the treatment of cystic fibrosis (CF):

- in patients aged 6 years and older and weighing 25 kg or more who have one of the following mutations in their cystic fibrosis transmembrane conductance regulator (*CFTR*) gene: *G551D*, *G1244E*, *G1349D*, *G178R*, *G551S*, *S1251N*, *S1255P*, *S549N*, *S549R* or *R117H*.

KALYDECO granules (13.4 mg, 25 mg, 50 mg, and 75 mg) are for the treatment of children with cystic fibrosis (CF):

- in patients aged 2 months and older and weighing 3 kg to less than 25 kg who have one of the following mutations in their *CFTR* gene: *G551D*, *G1244E*, *G1349D*, *G178R*, *G551S*, *S1251N*, *S1255P*, *S549N*, *S549R* or *R117H*.

KALYDECO is not for use in people with CF with two copies of the *F508del* mutation.

KALYDECO is not approved in patients under 2 months of age or weighing less than 3 kg.

How does KALYDECO work?

KALYDECO belongs to a group of medicines called “*cystic fibrosis transmembrane conductance regulator (CFTR) potentiators*.” The *CFTR* protein is a channel at the surface of the cell that allows the movement of particles such as chloride in and out of the cell, contributing to salt and water balance. KALYDECO helps this *CFTR* protein channel open more often allowing more chloride to pass through.

What are the ingredients in KALYDECO?

Medicinal ingredient: ivacaftor.

Non-medicinal ingredients:

- **Tablets** (150 mg): carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose acetate succinate, indigo carmine aluminum lake, lactose monohydrate, magnesium stearate, microcrystalline cellulose, PEG 3350, polyvinyl alcohol, sodium lauryl sulfate, talc, and titanium dioxide. The printing ink contains ammonium hydroxide, iron oxide black, propylene glycol, and shellac.
- **Granules** (13.4 mg, 25 mg, 50 mg, and 75 mg): colloidal silicon dioxide, croscarmellose sodium, hypromellose acetate succinate, lactose monohydrate, magnesium stearate, mannitol, sucralose, and sodium lauryl sulfate.

KALYDECO comes in the following dosage forms:

- **Tablets:** 150 mg.
- **Granules:** 13.4 mg, 25 mg, 50 mg, and 75 mg.

Do not use KALYDECO if:

- you are allergic (hypersensitive) to ivacaftor or to any of the non-medicinal ingredients or any of the ingredients in the component of the container.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take KALYDECO. Talk about any health conditions or problems you may have, including if you:

- have kidney or liver disease.
- have a problem with the lens of your eye (cataract). Your doctor may recommend eye exams before and after your treatment with KALYDECO.
- are pregnant or plan to become pregnant. It is not known if KALYDECO will harm your unborn baby. You and your doctor should decide if you will take KALYDECO while you are pregnant.
- are breastfeeding or planning to breastfeed. It is not known if KALYDECO passes into your breast milk. You and your doctor should decide if you will take KALYDECO while you are breastfeeding.

Other warnings you should know about:

Driving and using machines: You may get dizzy when you take KALYDECO. Wait to see how you feel after taking KALYDECO before you drive or use machines. **For children:** you or your child's caregiver should supervise your child when they take KALYDECO. Wait to see if your child is dizzy after taking KALYDECO before they ride their bikes or do anything else that needs their full attention.

Abnormal blood test results: KALYDECO can cause abnormal blood test results. Your doctor will decide if blood tests are needed.

If you have had high liver enzymes in the past, your doctor may order blood tests to check your liver more often.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with KALYDECO:

- antifungal medicines, such as ketoconazole, itraconazole, posaconazole, voriconazole, fluconazole.
- antibiotics used for the treatment of bacterial infection, such as clarithromycin, erythromycin, rifampin, rifabutin.
- phenobarbital, carbamazepine, phenytoin used for the treatment of epileptic seizures
- oral midazolam, alprazolam, diazepam, triazolam used for the treatment of anxiety, insomnia, agitation, etc.
- digoxin is used for the treatment of mild to moderate congestive heart failure and an abnormal heart rhythm called atrial fibrillation.
- cyclosporine, tacrolimus are used after an organ transplantation.
- warfarin is an anticoagulant used to prevent heart attacks, stroke, and blood clots.
- St. John's wort (*Hypericum perforatum*) is an herbal medicine.
- grapefruit juice and grapefruit.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How to take KALYDECO:

- Always take KALYDECO exactly how your doctor tells you.
- Take your doses of KALYDECO 12 hours apart.
- Even if you feel well, keep taking KALYDECO exactly how your doctor tells you.
- Check with your doctor if you are not sure about your dose.
- Do not change the dose or stop taking the medicine without first talking to your doctor.
- Your doctor may need to adjust your dose if you have liver disease or if you are taking medications that may interact with KALYDECO.

How to prepare the granules:

- Each KALYDECO box contains 4 individual wallets.
- There are 7 days of granules in each wallet. Each wallet contains 14 packets of granules (7 morning doses and 7 evening doses).
- Finish all doses from one wallet before starting a new one.
- **Morning Dose:** Remove the first dose from the wallet.
- Hold the packet with cut line on top.
- Shake the packet gently to settle the KALYDECO granules.
- Tear or cut packet open along cut line.
- Carefully pour all of the KALYDECO granules in the packet into 1 teaspoon of soft food or liquid. Food or liquid should be at or below room temperature. Each packet is for single use only. Some examples of soft foods or liquids include puréed fruits or vegetables, yogurt, applesauce, water, milk, breast milk, infant formula, or juice (except for grapefruit juice).
- Mix the KALYDECO granules with food or liquid.
- **After mixing, give KALYDECO within 1 hour. Make sure all medicine is taken. This is very important for KALYDECO to work properly and be effective.**
- If mixed with food, KALYDECO granules must not be stored for future use.
- **Evening Dose:** This dose should be taken 12 hours after your **Morning Dose**. Follow the same steps listed above under **Morning Dose**.

KALYDECO should always be taken with a fat containing food:

- Taking KALYDECO with fat containing food is important to get the right amount of medicine in your body.
- Each dose should be taken just before or just after fat containing food.
- Meals and snacks recommended in CF guidelines or meals recommended in standard nutritional guidelines contain adequate amounts of fat. A serving size of foods appropriate for age from a typical CF diet should be given.
- Examples of meals that contain fat:
 - meals prepared with butter or oils.
 - meals that have eggs, nuts, whole milk dairy products (such as whole milk, cheese, and yogurt) or meats.

Usual dose:

KALYDECO Tablets (for patients aged 6 years and older and weighing 25 kg or more):

One tablet every 12 hours by mouth with fat containing food (see below). Do not crush or chew the tablet. Swallow the tablet whole.

- Each KALYDECO box contains 4 individual blister cards.
- There are 7 days of tablets for each blister card. Each blister card contains 14 tablets (7 morning doses and 7 evening doses).

- You may cut along the dotted line to separate your doses from the blister card.
- **Morning Dose:** unpeel the paper backing from a blister card. Do not push the tablet through the paper backing because the tablets could break. Remove one KALYDECO tablet and swallow it whole with food that contains fat.
- **Evening Dose:** 12 hours after your first dose, unpeel the paper backing from another blister card to remove one KALYDECO tablet and swallow it whole with food that contains fat.

KALYDECO Granules (for children aged 2 months and older and weighing 3 kg to less than 25 kg): Your doctor will prescribe the appropriate amount based on age and weight. The usual dose is to take 1 packet of granules every 12 hours with fat-containing food.

Overdose:

If you think you, or a person you are caring for, have taken too much KALYDECO, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

- If you miss a dose of KALYDECO and it is **within 6 hours** of when you usually take it, take your missed dose of KALYDECO as prescribed with fat-containing food as soon as possible.
- If you miss a dose of KALYDECO and it is more **than 6 hours** after the time you usually take it, **skip** that dose only and take the next dose when you usually take it. Do not take 2 doses at the same time to make up for your missed dose.

What are possible side effects from using KALYDECO?

These are not all the possible side effects you may have when taking KALYDECO. If you experience any side effects not listed here, tell your healthcare professional.

Side effects of KALYDECO may include:

- | | |
|--|--|
| <ul style="list-style-type: none"> • Diarrhea • Nausea • Stomach (abdominal) pain • Common cold • Runny nose • Upper respiratory tract infection • Changes in the type of bacteria in your sputum | <ul style="list-style-type: none"> • Headache • Dizziness • Nasal congestion • Sinus congestion • Sore throat • Rash • Joint pain |
|--|--|

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Abdominal pain (stomach)	√		

Liver problems: Yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite, pale stools, itchy skin		√	
Low blood sugar (glucose): Sweating, shakiness, weakness, hunger, nausea, dizziness, headache	√		
UNKNOWN FREQUENCY			
Allergic reaction: rash; swelling of the face, lips, tongue or throat; difficulty swallowing or breathing			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- **Tablets:** Store at 20-25°C; short periods at 15-30°C are permitted.
- **Granules:** Store at or below 30°C.

Keep out of reach and sight of children.

If you want more information about KALYDECO:

- Talk to your healthcare professional.
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website (<http://www.vrtx.ca>), or by calling 1-877-634-8789.

This leaflet was prepared by Vertex Pharmaceuticals (Canada) Incorporated.

Last Revised: Nov 27,2023

© 2023 Vertex Pharmaceuticals (Canada) Incorporated

Vertex and KALYDECO are registered trademarks of Vertex Pharmaceuticals Incorporated.